



ELSEVIER

Available online at www.sciencedirect.com

**LINEAR ALGEBRA
AND ITS
APPLICATIONS**

Linear Algebra and its Applications 368 (2003) 379–385

www.elsevier.com/locate/laa

On the Laplacian spectral radius of a tree

Ji-Ming Guo

Department of Mathematics, University of Petroleum, ShanDong 257061, China

Received 1 March 2001; accepted 27 November 2002

Submitted by R.A. Brualdi

Abstract

Let G be a graph; its Laplacian matrix is the difference of the diagonal matrix of its vertex degrees and its adjacency matrix. In this paper, we present a sharp upper bound for the Laplacian spectral radius of a tree in terms of the matching number and number of vertices, and deduce from that the largest few Laplacian spectral radii over the class of trees on a given number of vertices.

© 2003 Published by Elsevier Science Inc.

AMS classification: 05C50

Keywords: Laplacian spectral radius; Matching number; Retraction

1. Introduction

Let $G = (V, E)$ be a graph on vertex set $V = \{v_1, v_2, \dots, v_n\}$. Let $|G|$ denote the number of vertices of G . The Laplacian matrix $L(G) = D(G) - A(G)$ is the difference of $D(G) = \text{diag}(d(v_1), d(v_2), \dots, d(v_n))$, the diagonal matrix of vertex degrees, and the adjacency matrix. The Laplacian spectral radius of G is the largest eigenvalue of its Laplacian matrix, denoted by $\lambda_1(G)$. Since the 1980s, many authors, for example Anderson and Morley [1], Li and Zhang [5,6], Merris [7], and Rojo et al. [9] have done a lot of work on the Laplacian spectral radius of graphs. In this paper, a sharp upper bound for the Laplacian spectral radius of a tree is given in terms of the matching number and number of vertices.

Two distinct edges in a graph G are independent if they are not incident with a common vertex in G . A set of pairwise independent edges of G is called a matching

E-mail address: gjm2248@sina.com (J.-M. Guo).

in G , while a matching of maximum cardinality is a maximum matching in G . The matching number $\beta(G)$ of G is the cardinality of a maximum matching of G .

Throughout this paper, we shall denote the characteristic polynomial of $L(G)$ by $\mu(G, x) = \det(xI - L(G))$.

2. Lemmas and results

Lemma 1 [2]. *The maximum eigenvalue r' of every principal submatrix (of order less than n) of a non-negative matrix A (of order n) does not exceed the maximum eigenvalue r of A . If A is irreducible, then $r' < r$ always holds. If A is reducible, then $r' = r$ holds for at least one principal submatrix.*

Lemma 2 [2]. *The increase of any element of a non-negative matrix A does not decrease the maximum eigenvalue. The maximum eigenvalue increases strictly if A is an irreducible matrix.*

Lemma 3 [3]. *Let G be a bipartite graph. Then $B(G) = D(G) + A(G)$ and $L(G)$ are unitarily similar; in particular, the maximum eigenvalue of $L(G)$ is simple provided G is connected.*

From Lemmas 1–3, we have the following result:

Corollary 1. *Let G be a connected bipartite graph, and let G' be a subgraph of G . Then $\lambda_1(G') \leq \lambda_1(G)$, and equality holds if and only if $G' = G$.*

Lemma 4 [3]. *Let T be a tree. Suppose v_1 and v_k are vertices each of degree at least three. Suppose the unique path v_1, v_2, \dots, v_k from v_1 to v_k is homeomorphic to an edge (i.e., apart from the endpoints, each vertex on the path has degree two). Let T' be the tree obtained from T by retracting along the entire path (i.e., deleting all edges $\{v_1v_2, v_2v_3, \dots, v_{k-1}v_k\}$ and identifying vertices v_1, v_2, \dots, v_k). Then $\lambda_1(T') > \lambda_1(T)$.*

Lemma 5 [1]. *Let G be a graph. Then $\lambda_1(G) \leq \max\{d(u) + d(v) : uv \in E\}$.*

Lemma 6 [8]. *If G has at least one edge, then $\lambda_1(G) \geq d_1(G) + 1$. For G a connected graph on $n > 1$ vertices, equality holds if and only if $d_1(G) = n - 1$.*

Let T_m^n ($2m \leq n + 1$) denote the tree graph with n vertices obtained from a star $K_{1,n-m}$ by joining $m - 1$ pendants (degree 1 vertices) of $K_{1,n-m}$ to the $m - 1$ isolated vertices by $m - 1$ edges.

Theorem 1. Let T be a tree on n vertices with matching number $\beta = \beta(T)$. Then $\lambda_1(T) \leq r$, where r is the maximum root of the equation

$$x^3 - (n - \beta + 4)x^2 + (3n - 3\beta + 4)x - n = 0.$$

The equality holds if and only if $T = T_\beta^n$.

Proof. By the direct computation, we have the characteristic polynomial of the tree T_β^n is

$$\begin{aligned} \mu(T_\beta^n, x) &= x(x-1)^{n-2\beta}(x^2-3x+1)^{\beta-2} \\ &\quad \times [x^3 - (n - \beta + 4)x^2 + (3n - 3\beta + 4)x - n]. \end{aligned}$$

Suppose that v_1 and v_2 are vertices of T each of degree at least three and the unique path from v_1 to v_2 is homeomorphic to an edge. Then we can apply Lemma 4 to T . Thus, a new tree T_1 can be constructed by retracting T along the entire path. Go on the above process until we obtain a tree $T_i = \tilde{T}$ ($T_0 = T$) such that we can not apply Lemma 4 to \tilde{T} . Then, we have

$$|\tilde{T}| \leq |T|, \quad \beta(\tilde{T}) \leq \beta(T)$$

and

$$\lambda_1(T) \leq \lambda_1(\tilde{T}) \quad (\lambda_1(T) = \lambda_1(\tilde{T}) \text{ if and only if } \tilde{T} = T).$$

Further, we have $\beta(T) - \beta(\tilde{T}) \leq |T| - |\tilde{T}|$ and there exists at most one vertex w of \tilde{T} such that $d(w) \geq 3$. If there is no such vertex w , then \tilde{T} is a path. From the above discussions, we conclude that $T = \tilde{T} = P_n$, a path with n vertices. Since $\lambda_1(P_n) = 4 \sin^2((n-1)\pi/2n)$ (see [1]) and $\beta(P_n) = \lfloor n/2 \rfloor$, the result is obvious. If there is such a vertex w such that $d(w) \geq 3$. We distinguish the following two cases:

Case 1. Assume that $\tilde{T} = K_{1,|\tilde{T}|-1}$ or $T_{\beta(\tilde{T})}^{|\tilde{T}|}$. Since $\beta(T) - \beta(\tilde{T}) \leq |T| - |\tilde{T}|$, then \tilde{T} is an induced subgraph of T_β^n . We have from Corollary 1 that $\lambda_1(\tilde{T}) \leq \lambda_1(T_{\beta(T)}^n)$, the equality holds if and only if $T = \tilde{T} = K_{1,n-1}$ or $T_{\beta(T)}^n$.

Case 2. Assume that $\tilde{T} \neq K_{1,|\tilde{T}|-1}$, $T_{\beta(\tilde{T})}^{|\tilde{T}|}$. Then there exists a $w-v$ path of \tilde{T}

$$W : w = w_0, w_1, w_2, \dots, w_{k-1}, w_k = v,$$

such that $k \geq 3$ and v is a pendant vertex of \tilde{T} .

Let $T^* = \tilde{T} - w_{k-2}w_{k-1} + ww_{k-1}$, then $|T^*| = |\tilde{T}|$. Further, we verify that $\beta(T^*) = \beta(\tilde{T})$. Suppose M is a maximum matching of \tilde{T} . Then either $w_{k-2}w_{k-1} \in M$ or $w_{k-1}w_k \in M$. If $w_{k-2}w_{k-1} \in M$, then $M^* = \{M - \{w_{k-2}w_{k-1}\}\} \cup \{w_{k-1}w_k\}$ is a matching of T^* and $|M^*| = |M|$; if $w_{k-2}w_{k-1} \notin M$, then $w_{k-1}w_k \in M$. Thus, M is also a matching of T^* . So, we have $\beta(T^*) \geq |M| = \beta(\tilde{T})$. By similar reasoning, we have $\beta(T^*) \leq \beta(\tilde{T})$. Hence, we have $\beta(T^*) = \beta(\tilde{T})$.

We have from Lemmas 5 and 6 that

$$\lambda_1(T^*) > d_1(T^*) + 1 = d_1(\tilde{T}) + 2 \geq \lambda_1(\tilde{T}).$$

Continuing the above process we have $\lambda_1(\tilde{T}) < \lambda_1(T_{\beta(\tilde{T})}^{|\tilde{T}|})$. From the discussion of Case 1, We conclude that $\lambda_1(T) < \lambda_1(T_{\beta(T)}^n)$, and the proof is complete. \square

We have from Lemmas 4 and 5 that

$$\lambda_1(T_i^n) > d_1(T_i^n) + 1 = n - i + 1 = d_1(T_{i+1}^n) + 2 \geq \lambda_1(T_{i+1}^n). \quad (1)$$

Then, the following two results hold.

Corollary 2 [1]. *Let T be a tree of order n , then $\lambda_1(T) \leq n$, and equality holds if and only if $T = T_1^n = K_{1,n-1}$.*

Corollary 3. *Let T be a tree of order n . If $T \neq K_{1,n-1}$ then $\lambda_1(T) \leq r_1$, where r_1 is the maximum root of the equation*

$$x^3 - (n+2)x^2 + (3n-2)x - n = 0.$$

The equality holds if and only if $T = T_2^n$.

Let $T(s, t)$ denote the tree of diameter 3 having exactly two non-pendant vertices, one of which is adjacent to s pendants and the other of which is adjacent to t pendants. In particular $n = s + t + 2$. Let

$$F = \{T(s, t) \mid s + t + 2 = n, s \geq 1, t \geq 1\}.$$

Corollary 4. *Let T be a tree with $n \geq 6$ vertices, $T \neq K_{1,n-1}$ and $T \neq T(n-3, 1) = T_2^n$. Then $\lambda_1(T) \leq r_2$, where r_2 is the maximum root of the equation*

$$x^3 - (n+2)x^2 + (4n-7)x - n = 0.$$

The equality holds if and only if $T = T(n-4, 2)$.

Proof. We have from [4] that the characteristic polynomials of $L(T(s, t))$ and $L(T(s+1, t-1))$ are

$$\mu(T(s, t), x) = x(x-1)^{n-4}[x^3 - (n+2)x^2 + (2n+st+1)x - n],$$

$$\begin{aligned} \mu(T(s+1, t-1), x) &= x(x-1)^{n-4}[x^3 - (n+2)x^2 \\ &\quad + (2n+(s+1)(t-1)+1)x - n]. \end{aligned}$$

In particular,

$$\mu(T(n-4, 2), x) = x(x-1)^{n-4}[x^3 - (n+2)x^2 + (4n-7)x - n].$$

and

$$\mu(T(n-5, 3), x) = x(x-1)^{n-4}[x^3 - (n+2)x^2 + (5n-14)x - n].$$

Let

$$f_1(x) = x^3 - (n+2)x^2 + (2n+st+1)x - n,$$

$$f_2(x) = x^3 - (n+2)x^2 + (2n+(s+1)(t-1)+1)x - n,$$

$$f(x) = x^3 - (n+2)x^2 + (4n-7)x - n,$$

$$g(x) = x^3 - (n+2)x^2 + (5n-14)x - n.$$

If $s \geq t, x > 0$, then $f_1(x) - f_2(x) = (s-t+1)x > 0$, we have

$$\lambda_1(T(s, t)) < \lambda_1(T(s+1, t-1)) \quad (s \geq t). \quad (2)$$

Let $H(s, t)$ denote trees of diameter 4 obtained from $T(s, t)$ by removing the non-pendant edge $e = uv$ and adding a new vertex w and edges uw and vw . In particular $n = s + t + 3$. Suppose $s \geq t \geq 1$. Let $H = \{H(s, t) \mid s + t + 3 = n, s \geq 1, t \geq 1\}$, then we have from Lemmas 4 and 5 that

$$\begin{aligned} \lambda_1(H(s, t)) &\leq d_1(H(s, t)) + 2 = s + 3 \\ &= d_1(H(s+1, t-1)) + 1 < \lambda_1(H(s+1, t-1)). \end{aligned} \quad (3)$$

By the direct computation, we have

$$\begin{aligned} \mu(H(n-4, 1), x) &= x(x-1)^{n-5} \\ &\quad \times [x^4 - (n+3)x^3 + (5n-4)x^2 - (6n-10)x + n]. \end{aligned}$$

$$\mu(T_3^n, x) = x(x-1)^{n-6}(x^2-3x+1)[x^3 - (n+1)x^2 + (3n-5)x - n].$$

Let

$$h(x) = x^4 - (n+3)x^3 + (5n-4)x^2 - (6n-10)x + n,$$

$$i(x) = x^3 - (n+1)x^2 + (3n-5)x - n.$$

Then

$$(x-2)i(x) - h(x) = x^2 - nx + n.$$

Note that when $h(x) = 0$, we have

$$(x^2 - 2x - 1)(x^2 - nx + n) = (x-2)[x^2 - (2n-5)x + n].$$

The right-hand side is easily seen to be negative when $4 \leq n-2 \leq x \leq n-1$ and since

$$n-2 \leq \lambda_1(H(n-4, 1)) \leq n-1.$$

We see that at $x = \lambda_1(H(n-4, 1))$, we have $i(x) < 0$. Since $i(n-1) > 0$, it now follows that $\lambda_1(T_3^n) > \lambda_1(H(n-4, 1))$. Since

$$i(x) - f(x) = x^2 - (n-2)x > 0, \quad (x > n-2),$$

we have $\lambda_1(T_3^n) < \lambda_1(T(n-4, 2))$. Clearly, if $\beta(T) = 2$, then either $T \in F$ or $T \in H$. From (2) and (3), we conclude that if T is a tree on n vertices with matching number two and $T \neq T(n-3, 1)$, then

$$\lambda_1(T) \leq \max\{\lambda_1(T(n-4, 2)), \lambda_1(H(n-4, 1))\}.$$

From (1) and Theorem 1, we conclude that if T is a tree on n vertices with matching number $\beta(T) \geq 3$, then

$$\lambda_1(T) \leq \lambda_1(T_{\beta(T)}^n) \leq \lambda_1(T_3^n).$$

Since

$$\lambda_1(H(n-4, 1)) < \lambda_1(T_3^n) < \lambda_1(T(n-4, 2)).$$

Then, we complete this proof. \square

Corollary 5. Let T be a tree with $n \geq 6$ vertices, and let $T \neq K_{1,n-1}$, $T \neq T(n-3, 1)$ and $T \neq T(n-4, 2)$. Then $\lambda_1(T) \leq r_3$, where r_3 is the maximum positive root of the equation $x^3 - (n+1)x^2 + (3n-5)x - n = 0$. The equality holds if and only if $T = T_3^n$.

Proof. If $n = 6, 7$, then it is easy to show this result holds.

Suppose that $n \geq 8$. Since $T \neq K_{1,n-1}$, we have $\beta(T) \geq 2$. If $\beta(T) = 2$ and $T \neq T(n-3, 1)$, $T(n-4, 2)$, we have from (2) and (3) that

$$\lambda_1(T) \leq \max\{\lambda_1(T(n-5, 3)), \lambda_1(H(n-4, 1))\}.$$

Since for $n \geq 8$, $x < n-1$ ($x \in R$), we have $g(x) - i(x) = -x^2 + (2n-9)x > 0$. Then, for $n \geq 8$, $\lambda_1(T_3^n) > \lambda_1(T(n-5, 3))$. Combined with $\lambda_1(T_3^n) > \lambda_1(H(n-4, 1))$. We have $\lambda_1(T) < \lambda_1(T_3^n)$.

From (1) and Theorem 1, we conclude that if T is a tree on n vertices with matching number $\beta(T) \geq 3$, then

$$\lambda_1(T) \leq \lambda_1(T_{\beta(T)}^n) \leq \lambda_1(T_3^n),$$

and equality holds if and only if $T = T_3^n$. We complete this proof. \square

In particular, we have

Corollary 6. Let T be a tree on $n = 2k$ vertices with a perfect matching. Then

$$\lambda_1(T) \leq \frac{k+2+\sqrt{k^2+4}}{2},$$

and equality holds if and only if $T = T_k^n$.

Proof. By the direct computation, we have the characteristic polynomial of $L(T_k^n)$ is

$$\begin{aligned}\mu(T_k^n, x) &= x(x^2 - 3x + 1)^{k-2}[x^3 - (k+4)x^2 + (3k+4)x - 2k] \\ &= x(x-2)(x^2 - 3x + 1)^{k-2}[x^2 - (k+2)x + k].\end{aligned}$$

By Theorem 1, we complete this proof. \square

Acknowledgement

The author is grateful to the anonymous referee for giving valuable suggestions towards improving the proofs of the original version of this paper.

References

- [1] W.N. Anderson, T.D. Morley, Eigenvalues of the Laplacian of a graph, *Linear and Multilinear Algebra* 18 (1985) 141–145.
- [2] D.M. Cvetkovic, M. Doob, H. Sachs, *Spectra of graph-theory and applications*, VEB Deutscher Verlag d. Wiss. Berlin, 1979; Academic Press, New York, 1979.
- [3] R. Grone, R. Merris, V.S. Sunder, The Laplacian spectrum of a graph, *SIAM J. Matrix Anal. Appl.* 11 (2) (1990) 218–238.
- [4] R. Grone, R. Merris, Ordering trees by algebraic connectivity, *Graphs Combin.* 6 (1990) 229–237.
- [5] J.-S. Li, X.-D. Zhang, A new upper bound for eigenvalues of the Laplacian matrix of a graph, *Linear Algebra Appl.* 265 (1997) 93–100.
- [6] J.-S. Li, X.-D. Zhang, On the Laplacian eigenvalues of a graph, *Linear Algebra Appl.* 285 (1998) 305–307.
- [7] R. Merris, A note on Laplacian graph eigenvalues, *Linear Algebra Appl.* 285 (1998) 33–35.
- [8] R. Merris, Laplacian matrices of graphs: a survey, *Linear Algebra Appl.* 197&198 (1994) 143–176.
- [9] O. Rojo, R. Soto, H. Rojo, An always non-trivial upper bound for Laplacian graph eigenvalues, *Linear Algebra Appl.* 312 (2000) 155–159.